

P.O. Box 8522 Portland, Maine 04104 207-856-6620 207-856-6864 (fax) Immunochemicals for
Infectious Disease
Research
www.ViroStat-Inc.com

RabMAB®

no stabilizing proteins and should be stored at 2-8°C until ready for use.

product information for

## Streptococcus pneumoniae CWPS

## I. Monoclonal Antibody (Rabbit) Specificity

Product No.'s Ig class

O494 IgG1 Specific to the common cell wall polysaccharide (urinary antigen) of S. pneumoniae. ELISA, IFA & LF appl.\*

O495 IgG1 Specific to the common cell wall polysaccharide (urinary antigen) of S. pneumoniae. ELISA, IFA & LF appl.\*

**II. Purified Preparations** 

Product No.'s

0494 0495

RabMAB® purified preparations consist of >90% pure rabbit monoclonal antibody which has been purified from culture medium by protein A chromatography or sequential differential precipitations. The final preparation is formulated to a protein concentration of 100 µg/ml in 0.01 M phosphate buffered saline, pH 7.2 and contains 0.1% sodium azide. Each vial contains 1.0 ml. This product contains

Working dilution must be determined by the user. Suggested starting ranges are 1:10-1:50 for IFA and 1:20-1:200 for ELISA.

## THESE PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

## III. Fluorescein Conjugates

Product No.'s

These MONOTOPE™ products consist of purified monoclonal antibody conjugated with high purity isomer I of fluorescein isothiocyanate. Care is taken to ensure complete removal of any free fluorescein from the final product. The final preparation is formulated to an antibody concentration of of 100 µg/ml in 0.01 M phosphate buffered saline, pH 7.2 containing 0.1% sodium azide plus bovine serum albumin at 10 mg/ml. Each vial contains 1.0 ml. This product should be stored at 2-8°C until ready for use. Avoid repeated freeze-thawing by storing multiple aliquots at -20°C. Applications for these products include direct FA staining of target antigen in a permissive tissue culture system. Working strength must be determined by the user for each specific application but a starting range of 1:5 - 1:20 is recommended. Acetone fixation of the antigen source is recommended prior to staining.

THESE PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Comments: \*0494 as conjugate with 0495 as capture